

On the

Number

of Real Zeros

of
Random Sparse Polynomial Systems

or...

how few real zeros
does a random fewnomial system have?



Josué Tonelli-Cueto
(Johns Hopkins University)

Joint work with...



Alperen A. ERGÜR



Maté L. TELEK

Both pictures taken in San Antonio, Texas, USA

How many zeros does it have?

$$\begin{cases} a_1 + b_1 X + \gamma_1 Y + c_1 XYZ^d = 0 \\ a_2 + b_2 X + \gamma_2 Y + c_2 XYZ^d = 0 \\ a_3 + b_3 X + \gamma_3 Y + c_3 XYZ^d = 0 \end{cases}$$

generically... d in \mathbb{C}^3 (intersection theory)

$$\leq 2 \text{ in } \mathbb{R}^3$$

$$\leq 1 \text{ in } \mathbb{R}_+^3$$

Kushnirenko's Question:

$$A_1, \dots, A_n \subseteq \mathbb{N}^n \quad t_i := \#A_i$$

$$f := \begin{cases} f_1 := \sum_{\alpha \in A_1} f_{1,\alpha} X^\alpha \\ \dots \\ f_n := \sum_{\alpha \in A_n} f_{n,\alpha} X^\alpha \end{cases}$$

What do we call
such a system
with few terms?

fewnomial

$$\mathcal{Z}_r(f, \mathbb{R}_+^n) := \{g \in \mathbb{R}_+^n \mid f(g) = 0, \det D_g f \neq 0\}$$

Is there a bound of the form

$$\# \mathcal{Z}_r(f, \mathbb{R}_+^n) \leq U(t_1, \dots, t_n, n)?$$


Kushnirenko's Question:

Propagandistic version

Do fewnomial systems
have few positive zeros?

polynomial \rightarrow МНОГОЧЛЕН \rightarrow МАЛОЧЛЕН \rightarrow fewnomial
 $\underbrace{\hspace{10em}}_{\text{many nomials}} \quad \underbrace{\hspace{10em}}_{\text{few nomials}} \quad \text{pocosnomio (es)}$
 $\hspace{10em} \text{gutxinomio (ea)}$

Khovanskii's Answer & Refinements



$$A \subseteq \mathbb{N}^n \quad t = \# A (\leq t_1 + \dots + t_n)$$

$$f := \{ f_i := \sum_{\alpha \in A} f_{i,\alpha} X^\alpha \quad (i=1, \dots, n)$$

(Khovanskii, 1991)

$$\# \mathcal{Z}_r(f, \mathbb{R}_+^n) \leq 2^{\binom{t-1}{2}} (n+1)^{t-1}$$

(Bihan & Sottile; 2007)

$$\# \mathcal{Z}_r(f, \mathbb{R}_+^n) \leq \frac{e^2 + 3}{4} 2^{\binom{t-n-1}{2}} n^{t-n-1}$$

Kushnirenko's Question:

A more precise version

$$A_1, \dots, A_n \subseteq \mathbb{N}^n \quad t_i := \#A_i$$

$$f := \begin{cases} f_1 := \sum_{\alpha \in A_1} f_{1,\alpha} X^\alpha \\ \dots \\ f_n := \sum_{\alpha \in A_n} f_{n,\alpha} X^\alpha \end{cases}$$

! STILL
OPEN
FOR $n=2!!!$
...

$$\mathcal{Z}_r(f, \mathbb{R}_+^n) := \{g \in \mathbb{R}_+^n \mid f(g) = 0, \det D_g f \neq 0\}$$

Is there a bound of the form

$$\#\mathcal{Z}_r(f, \mathbb{R}_+^n) \leq \text{poly}(t_1, \dots, t_n)^n ?$$

Can we say

SOMETHING

for a random fewnomial system?

spoiler: yes!

Reasons to care...

about random fewnomial systems

Better understanding

of typical behaviour

— not the worst case, but the average typical case

Robust probabilistic results

might be a pathway to deterministic ones

$$\sup_{\mathcal{F}} \# \mathcal{Z}_r(\mathcal{F}, \mathbb{R}_+^n) = \sup_{e \geq 1} \left(\mathbb{E}_{\mathcal{F}} \# \mathcal{Z}_r(\mathcal{F}, \mathbb{R}_+^n)^e \right)^{1/e}$$

Our Random Model

$A_1, \dots, A_n \subseteq \mathbb{R}^n$ w/ $t_i = \#A_i$; Fixed

$$F := \begin{cases} F_1 := \sum_{\alpha \in A_1} f_{1,\alpha} X^\alpha \\ \dots \\ F_n := \sum_{\alpha \in A_n} f_{n,\alpha} X^\alpha \end{cases}$$

random
polynomial
system

such that the $f_{i,\alpha}$ are independent,
centered ($\mathbb{E} f_{i,\alpha} = 0$)
& Gaussian

presented at MEGA2019!

(Bürgisser, Ergür & TC; 2019) & (Bürgisser, ISSAC'23)
are particular cases of our model

Our Result: The Easy Form

$A_1, \dots, A_n \subseteq \mathbb{R}^n$ w/ $t_i = \#A_i$ Fixed

$$F := \left\{ F_i := \sum_{\alpha \in A_i} f_{i,\alpha} X^\alpha \quad (i = 1, \dots, n) \right.$$

s.t. the $f_{i,\alpha}$ are independent, centered & Gaussian

$$\mathbb{E} \sum_{F} \chi_r(F, \mathbb{R}_+^n) \leq \frac{1}{4^n} \prod_{i=1}^n t_i (t_i - 1)$$

Our Result: Unit Variance Case

$A_1, \dots, A_n \subseteq \mathbb{R}^n$ w/ $t_i = \#A_i$ Fixed

$$F := \left\{ F_i := \sum_{\alpha \in A_i} f_{i,\alpha} X^\alpha \quad (i = 1, \dots, n) \right.$$

s.t. the $f_{i,\alpha}$ are independent, centered & Gaussian

IF (VM1) for all i and $\alpha \in A_i$, $V(f_{i,\alpha}) \leq 1$

(VM2) for all i and $\alpha \in A_i$ vertex of $P_i := \text{conv}(A_i)$,

$$V(f_{i,\alpha}) = 1$$

THEN

$$\mathbb{E} \sum_{F \in \mathcal{F}} \chi(F, \mathbb{R}_+^n) \leq \frac{1}{4^n} V\left(\sum_{i=1}^n P_i\right) \prod_{i=1}^n (t_i - 1)$$

↑ #vertices
↑ Minkowski sum

Improves (Bürgisser, ISSAC'23)

Our Result: The Unmixed Case

$A \subseteq \mathbb{R}^n$ w/ $t = \#A$ Fixed

$$F := \left\{ F_i := \sum_{\alpha \in A} f_{i,\alpha} X^\alpha \quad (i = 1, \dots, n) \right.$$

s.t. the $f_{i,\alpha}$ are independent, centered & Gaussian

IF $\left\{ \begin{array}{l} \text{(VM1) for all } i \text{ and } \alpha \in A, V(f_{i,\alpha}) \leq 1 \\ \text{(VM2) for all } i \text{ and } \alpha \in A \text{ vertex of } P := \text{conv}(A), V(f_{i,\alpha}) = 1 \end{array} \right.$

OR

$V(f_{i,\alpha})$ only depends on $\alpha \in A$

THEN

$$\mathbb{E} \sum_{F} \mathcal{Z}_\nu(F, \mathbb{R}_+^n) \leq \frac{n+1}{4^n} \binom{t}{n+1}$$

Improves (Bürgisser, Ergür & TC; 2019)


Regular (Mixed) Subdivisions?

$$A \subseteq \mathbb{R}^n$$

$\pi: A \rightarrow \mathbb{R}$ lifting

upper envelope of A wr π

$$\mathcal{L}(A, \pi) := \text{conv} \left\{ \begin{pmatrix} \pi(\alpha) - s \\ \alpha \end{pmatrix} \mid \alpha \in A, s \geq 0 \right\}$$



Our Result: In all its detail

$A_1, \dots, A_n \subseteq \mathbb{R}^n$ w/ $t_i = \#A_i$ fixed

$$F := \left\{ F_i := \sum_{\alpha \in A_i} f_{i,\alpha} X^\alpha \quad (i = 1, \dots, n) \right.$$

s.t. the $f_{i,\alpha}$ are independent, centered & Gaussian

$\mu_{F,i}: A_i \rightarrow \mathbb{R}$ given by $\mu_{F,i}(\alpha) = \frac{1}{2} \ln V(F_{i,\alpha})$

$$\mathbb{E} \sum_{F} \tilde{Z}_V(F, \mathbb{R}_+^n) \leq \frac{1}{4^n} V \left(\sum_{i=1}^n \mathcal{L}(A_i, \mu_{F,i}) \right) \prod_{i=1}^n (t_i - 1)$$

vertices # Minkowski sum

⚠ # vertices of regular mixed subdivision induced by variance on supports

Three Tools

that made this possible

Tool I: Kac-Rice Formula

under some technical assumptions...

$$\mathbb{E} \mathcal{Z}(f, \Omega) = \int_{\Omega} \mathbb{E}(|\det D_x f| | f(x) = 0) \delta_{f(x)}(0) dx$$

conditional expectation

density of $f(x)$

Tool II: Cauchy-Binet Formula

(aka the Fundamental Lemma of Fewnomial Theory)


$$A, B \in \mathbb{F}^{m \times n}$$

$$\det(A B^T) = \sum_{\substack{J \subseteq \{1, \dots, n\} \\ \# J = m}} \det(A_J) \det(B_J)$$

where $A_J = \left(A_{i,j} \right)_{\substack{i \in \{1, \dots, m\} \\ j \in J}}$, $B_J := \left(B_{i,j} \right)_{\substack{i \in \{1, \dots, m\} \\ j \in J}}$

used also in the work of Bihan & Sottile

Tool III: Normal Fan



Open Questions

What about higher moments?

In particular, what about

$$\bigvee_F \mathbb{Z}_\nu(F, \mathbb{R}_+^n) ?$$

What about more general
Gaussian fewnomial systems?

Vielen Dank

für

Aufmerksamkeit!